

PRODUCT MONOGRAPH

PrURSO®
250 mg Tablets

PrURSO DS®
500 mg Tablets

ursodiol tablets

USP

Bile Acid Preparation

A05AA02

APTALIS PHARMA CANADA INC.
597 Laurier Blvd.
Mont-St-Hilaire, Quebec
Canada J3H 6C4

Date of Preparation:
November 2, 2011

Date of Revision:
August 18, 2014

Submission Control Number:175562

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....3
SUMMARY PRODUCT INFORMATION.....3
INDICATIONS AND CLINICAL USE3
CONTRAINDICATIONS4
WARNINGS AND PRECAUTIONS4
ADVERSE REACTIONS.....5
DRUG INTERACTIONS.....10
DOSAGE AND ADMINISTRATION11
OVERDOSAGE12
ACTION AND CLINICAL PHARMACOLOGY12
STORAGE AND STABILITY.....13
SPECIAL HANDLING INSTRUCTIONS.....13
DOSAGE FORMS, COMPOSITION AND PACKAGING14

PART II: SCIENTIFIC INFORMATION15
PHARMACEUTICAL INFORMATION.....15
CLINICAL TRIALS.....16
DETAILED PHARMACOLOGY18
TOXICOLOGY.....19
REFERENCES.....22

PART III: CONSUMER INFORMATION.....26

**URSO®
URSO DS®**

ursodiol tablets

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Non-medicinal Ingredients
oral	tablet 250 mg, 500 mg	None <i>For a complete listing see DOSAGE FORMS, COMPOSITION AND PACKAGING section.</i>

INDICATIONS AND CLINICAL USE

URSO® and URSO DS® (ursodiol), also known as ursodeoxycholic acid (UDCA) are indicated for:

- the management of cholestatic liver diseases, such as primary biliary cirrhosis (PBC).

Cholestatic liver diseases are characterized by a decrease in bile secretion and bile flow. Caution has to be exercised to maintain the bile flow of the patients taking UDCA..

The diagnosis of cholestatic liver diseases is based on the biochemical signs of cholestasis (such as an increase in alkaline phosphatase, γ -GT, bilirubin), and also an increase in IgM levels and the presence of antimitochondrial antibodies in PBC.

The monitoring of URSO® and URSO DS® in the management of cholestatic liver diseases should be based on the biochemical parameters of cholestasis, as described above, as well as on signs of hepatic cytolysis (such as AST, ALT) which are very often associated with cholestasis during the progression of the diseases.

Serum liver function tests (γ -GT, alkaline phosphatase, AST, ALT) and bilirubin level should be monitored every month for three months after start of therapy, and every six months thereafter. Improved serum liver function tests (e.g. AST, ALT) do not always correlate with improved disease status. In addition to identifying responsive and non-responsive patients, this monitoring will allow the early detection of a possible deterioration of the hepatic function. For patients who have a recent history of adequate biochemical response to the treatment, UDCA discontinuation should be considered when serum liver function tests increase to a level considered clinically significant, generally increase in ALT, AST levels three times the baseline value and increase in total bilirubin to twice the baseline value, confirmed by repeated tests. (see WARNINGS and

PRECAUTIONS and DOSAGE and ADMINISTRATION).

URSO[®] and URSO DS[®] are not indicated for the treatment of decompensated cirrhosis.

Geriatrics:

Appropriate studies with URSO[®] and URSO DS[®] have not been performed in the geriatric population. However, geriatric-specific problems that would limit the use or usefulness of URSO[®] and URSO DS[®] in the elderly are not expected.

Pediatrics:

The safety and effectiveness of URSO[®] and URSO DS[®] in children have not been established.

CONTRAINDICATIONS

Patients with complete biliary obstruction of extrahepatic origin; patients with widespread intrahepatic obstruction and patients who are hypersensitive to ursodiol or to any ingredient in the formulation For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph.

WARNINGS AND PRECAUTIONS

Carcinogenesis and Mutagenesis

URSO[®] and URSO DS[®] have no carcinogenic, mutagenic or teratogenic effects in laboratory animals treated at higher doses than those intended for therapy in humans, and after long-term treatment (see TOXICOLOGY).

Hepatic/Biliary/Pancreatic

Patients with variceal bleeding, hepatic encephalopathy, ascites, or in need of an urgent liver transplant, should receive appropriate specific treatment. Caution should be exercised when UDCA is administered in a setting of partial biliary obstruction of extra-hepatic origin.

Special Populations

Pregnant Women: There are no adequate or well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, URSO[®] and URSO DS[®] should not be used in women who are or may become pregnant. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the foetus. (See also TOXICOLOGY.)

Nursing Women: It is not known whether ursodiol is excreted in human milk. Since many drugs are excreted in human milk, caution should be exercised when URSO[®] or URSO DS[®] is administered to a nursing mother.

Pediatrics: The safety and effectiveness of URSO[®] and URSO DS[®] in children have not been established.

Geriatrics: Appropriate studies with URSO[®] and URSO DS[®] have not been performed in the geriatric population. However, geriatric-specific problems that would limit the use or usefulness of URSO[®] and URSO DS[®] in the elderly are not expected.

Monitoring and Laboratory Tests

Lithocholic acid, one of the metabolites of ursodeoxycholic acid (URSO[®] and URSO DS[®]) is hepatotoxic unless it is effectively detoxified in the liver. Therefore, the following tests are important for patient monitoring:

Serum liver function tests (γ -GT, alkaline phosphatase, AST, ALT) and bilirubin levels should be monitored every month for three months after start of therapy, and every six months thereafter. Serial monitoring will allow for the early detection of a possible deterioration of the hepatic function. Serum levels of these parameters usually decrease rapidly. Improved serum liver tests (e.g. AST, ALT) do not always correlate with improved disease status. For patients who have a recent history of adequate biochemical response to the treatment, UDCA discontinuation should be considered when serum liver function tests increase to a level considered clinically significant (see DOSAGE and ADMINISTRATION and WARNINGS and PRECAUTIONS).

Caution has to be exercised to maintain the bile flow of the patients taking UDCA.

ADVERSE REACTIONS

Adverse Drug Reaction Overview

Adverse events observed in clinical trials are tabulated and described below. In a 180 patient placebo-controlled trial in primary biliary cirrhosis, the common adverse events (i.e. $\geq 1\%$) included leukopenia, skin rash, diarrhea, blood creatinine increased, blood glucose increased, and peptic ulcer. In a second trial with 60 patients, the frequency of treatment-emergent adverse event reporting was higher with the most common (defined as $\geq 5\%$) being asthenia, dyspepsia, edema peripheral, hypertension, nausea, GI disorder, chest pain, and pruritus. In this second trial there were 4 serious adverse events: 1 patient with diabetes mellitus, 1 patient with breast nodule and 2 patients with fibrocystic breast disease. None of these events were considered related to the medication. At the recommended dosage, ursodiol is well-tolerated and has no significant adverse events.

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

The adverse reactions in Table 1 below were observed in clinical trials in primary biliary cirrhosis with 180 patients (89 randomized to URSO[®] treatment, 91 to placebo treatment). Adverse events are reported regardless of attribution to the test medication. Adverse reactions occurring at a rate of 1% or higher in the URSO[®] group, and that are higher than placebo are included in Table 1.

Diarrhea and thrombocytopenia at 12 months, nausea/vomiting, fever and other side effects are not included, because they occurred at the same rate or a lower rate than placebo.

Table 1: Adverse events with a frequency \geq 1% Observed in a Clinical Trial of 180 patients

Adverse event (ordered by MedDRA System Organ Class)		Visit at 12 Months		Visit at 24 Months	
		UDCA ¹ n (%)	Placebo n (%)	UDCA ¹ n (%)	Placebo n (%)
Blood and lymphatic system disorders	Leukopenia	-	-	2 (2.63)	-
Gastrointestinal disorders	Diarrhea	-	-	1 (1.32)	-
	Peptic ulcer	-	-	1 (1.32)	-
Investigations	Blood creatinine increased	-	-	1 (1.32)	-
	Blood glucose increased	1 (1.18)	-	1 (1.32)	-
Skin and subcutaneous tissue disorders	Rash	-	-	2 (2.63)	-

¹ UDCA=Ursodeoxycholic acid=Ursodiol =URSO®

Note: Those AEs occurring at the same or higher incidence in the placebo as in the UDCA group have been deleted from this table (this includes diarrhea and thrombocytopenia at 12 months, nausea/vomiting, fever and other toxicity).

In a randomized, cross over study in sixty PBC patients, four patients experienced one serious adverse event each (diabetes mellitus, breast nodule, and fibrocystic breast disease (2 patients)). No deaths occurred in the study. Forty-three patients (43/71.7%) experienced at least one treatment-emergent adverse event (TEAEs) during the study. The most common (defined as \geq 5%) TEAEs were asthenia, (11.7%), dyspepsia (10%), edema peripheral (8.3%), hypertension (8.3%), nausea (8.3%), GI disorders (5%), chest pain (5%), and pruritus (5%). These nine TEAEs included abdominal pain and asthenia (1 patient), nausea (3 patients), dyspepsia (2 patients), and anorexia and esophagitis (1 patient each). One patient on the BID regimen (total dose 1000 mg) withdrew due to nausea. All of these nine TEAEs except esophagitis were observed with the BID regimen at a total daily dose of 1000 mg or greater.

Table 2: Treatment-Emergent Adverse Events (TEAEs) with a Frequency of \geq 1 % Observed in a Clinical Trial of 60 PBC patients

Adverse event (ordered by MedDRA System Organ Class)		TEAEs, n (%)
Blood and lymphatic system disorders	Anemia	1 (1.7)
	Lymphadenopathy	2 (3.3)
Cardiac disorders	Arrhythmia	2 (3.3)
	Cardiovascular disorder	2 (3.3)
Ear and labyrinth disorders	Deafness	1 (1.7)
	Vertigo	1 (1.7)
Eye disorders	Cataract	2 (3.3)
	Eye disorder	1 (1.7)
	Retinal disorder	1 (1.7)
Gastrointestinal disorders	Abdominal pain	2 (3.3)
	Diarrhea	2 (3.3)
	Dyspepsia	6 (10)

Adverse event (ordered by MedDRA System Organ Class)	TEAEs, n (%)	
	Dysphagia	1 (1.7)
	Esophagitis	1 (1.7)
	Flatulence	1 (1.7)
	Gastrointestinal disorder	3 (5.0)
	Nausea	5 (8.3)
	Salivary gland enlargement	1 (1.7)
	Stomach ulcer	1 (1.7)
General disorders and administration site conditions	Asthenia	7 (11.7)
	Chest pain	3 (5.0)
	Chest pain substernal	1 (1.7)
	Cyst	1 (1.7)
	Edema	5 (8.3)
	Edema generalized	1 (1.7)
	Edema peripheral	5 (8.3)
	Granuloma	1 (1.7)
	Hemorrhagic ulcer	1 (1.7)
	Pain	1 (1.7)
Hepatobiliary disorders	Biliary pain	1 (1.7)
Immune system disorders	Amyloidosis	1 (1.7)
Infections and infestations	Bronchitis	1 (1.7)
	Cystitis	1 (1.7)
	Herpes simplex	1 (1.7)
	Infection	1 (1.7)
	Otitis media	1 (1.7)
	Pharyngitis	1 (1.7)
	Pneumonia	1 (1.7)
	Rhinitis	2 (3.3)
	Urinary tract infection	1 (1.7)
	Vaginitis	1 (1.7)
Metabolism and nutrition disorders	Anorexia	1 (1.7)
	Diabetes mellitus	2 (3.3)
Musculoskeletal and connective tissue disorders	Back pain	1 (1.7)
	Bone disorder	1 (1.7)
	Bone fracture spontaneous	1 (1.7)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)	Breast neoplasm	1 (1.7)
	Lung nodule	1 (1.7)
	Plantar warts	1 (1.7)
Nervous system disorders	Dizziness	2 (3.3)
	Headache	1 (1.7)
	Migraine	1 (1.7)
	Paresthesia	1 (1.7)
Reproductive system and breast disorders	Breast nodule	1 (1.7)
	Fibrocystic breast disease	2 (3.3)
	Menorrhagia	1 (1.7)
Respiratory, thoracic and mediastinal disorders	Dyspnea	1 (1.7)
	Lung disorder	1 (1.7)
	Respiratory disorder	1 (1.7)
	Sore nose	2 (3.3)
Skin and subcutaneous tissue disorders	Acne	2 (3.3)
	Miliaria	1 (1.7)
	Pruritus	3 (5.0)

Adverse event (ordered by MedDRA System Organ Class)		TEAEs, n (%)
	Psoriasis	1 (1.7)
	Rash	1 (1.7)
	Skin disorder	2 (3.3)
	Skin hypertrophy	1 (1.7)
Vascular disorders	Hypertension	5 (8.3)

Less Common Clinical Trial Adverse Drug Reactions (<1%)

Analysis of the data in the trial with 180 patients (Table 1) revealed no reports of adverse events at rates <1 % with the exception of those adverse events that occurred at the same or at a higher incidence in the treatment group than placebo. No data for TEAEs occurring at rates <1 % in the trial of 60 patients (Table 2) are available due to the small sample size.

Abnormal Hematologic and Clinical Chemistry Findings

In the placebo-controlled trial with 180 patients, change from baseline in hematologic parameters and non-hepatic clinical chemistry were analyzed. Statistically significant differences from baseline are reported in Tables 3 and 4.

Table 3: Hematologic Parameters: Changes from Baseline

		Baseline		Endpoint		Change from Baseline	
		UDCA	Placebo	UDCA	Placebo	UDCA (± SD)	Placebo (± SD)
WBC	Mean (± SD)	5.9 (2.0)	6.2 (4.1)	5.5 (1.6)	5.8 (2.4)	-0.5** (1.4)	-0.5 (4.3)
	n	88	87	83	75		
Platelets	Mean (± SD)	238.5 (92.5)	245.4 (112.4)	211.2 (87.2)	223.9 (94.3)	-29.4** (39.3)	-17.7* (58.0)
	n	86	86	82	74		

* Statistically different from zero, $p < 0.05$

**Statistically different from zero, $p < 0.01$

There was a significant decrease ($p < 0.01$) in WBC and platelets in the UDCA-treated group from baseline and a significant ($p < 0.05$) decrease in platelets in the placebo group. There was no significant change in haemoglobin.

Table 4: Clinical Chemistries: Changes from Baseline

		Baseline		Endpoint		Change from Baseline	
		UDCA	Placebo	UDCA	Placebo	UDCA (± SD)	Placebo (± SD)
Calcium (mg/dL)	Mean (± SD)	9.49 ^a (0.40)	9.47 (0.40)	9.39 (0.43)	9.30 (0.51)	-0.12 ^{**a} (0.37)	-0.19 ^{**} (0.37)
	n	89	91	83	76		
Cholesterol (mg/dL)	Mean (± SD)	287.73 ^a (121.12)	276.03 (105.22)	223.53 (56.80)	261.46 (83.53)	-67.39 ^{**b} (93.31)	-11.32 [*] (47.70)
	n	89	91	83	76		
Creatinine (mg/dL)	Mean (± SD)	0.86 (0.19)	0.84 (0.21)	0.92 (0.19)	0.92 (0.26)	0.07 ^{**a} (0.18)	0.07 ^{**} (0.23)
	n	89	91	83	76		
Total Thyroxine (µg/dL)	Mean (± SD)	8.66 ^a (1.63)	8.60 (2.27)	7.96 (1.87)	8.27 (3.25)	-0.69 ^{**a} (1.52)	-0.49 (2.52)
	n	87	90	83	74		
Triglycerides (mg/dL)	Mean (± SD)	102.82 ^a (49.25)	117.11 (70.57)	114.18 (55.13)	121.52 (57.56)	11.76 ^{**a} (44.38)	3.00 (56.74)
	n	88	89	83	75		

** Statistically different from zero, $p < 0.01$

* Statistically different from zero, $p < 0.05$

^a $p = ns$, UDCA versus placebo

^b $p = 0.0001$, UDCA versus placebo

All the non-hepatic clinical chemistries at baseline were not significantly different ($p > 0.05$) between the UDCA- and placebo- treated groups. In the UDCA group there was a significant ($p > 0.05$) decrease from baseline in calcium, cholesterol and total thyroxine and a significant increase ($p > 0.05$) in creatinine and triglycerides. In the placebo group there was a significant ($p > 0.05$) decrease in cholesterol and significant increase ($p > 0.05$) in calcium and creatinine. There was no significant change seen for sodium, potassium, phosphorus, HDL, and AMA.

Post-Market Adverse Drug Reactions

The following adverse reactions, presented by system organ class in alphabetical order, have been identified during post-approval use of ursodiol. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- *Blood and lymphatic system disorders:* anemia, eosinophilia, leukopenia, neutropenia, thrombocytopenia.
- *Cardiac disorders:* palpitations.
- *Gastrointestinal disorders:* abdominal discomfort, abdominal pain, cheilitis, constipation, diarrhea, dyspepsia, nausea, vomiting.
- *General disorders and administration site conditions:* malaise, peripheral edema, pyrexia.
- *Hepatobiliary disorders:* jaundice (or aggravation of pre-existing jaundice)
- *Immune system disorders:* angioedema and laryngeal edema, drug hypersensitivity to include facial edema, urticaria.
- *Investigations:* blood glucose increased, blood urine present, weight decreased, weight increased, ALT increased, AST increased, blood alkaline phosphatase increased, blood bilirubin increased, γ -GT increased, transaminases increased. Rare instances of severe liver injury (elevated values for ALT/AST, ALP, γ -GTP and total bilirubin) have been reported with URSO.
- *Musculoskeletal and connective tissue disorders:* myalgia
- *Nervous system disorders:* dizziness, headache.
- *Respiratory, thoracic and mediastinal disorders:* cough, interstitial lung disease.
- *Skin and subcutaneous tissue disorder:* alopecia, dermatitis exfoliative, drug eruption, erythema, lichenoid keratosis, photosensitivity reaction, pruritus, rash.

DRUG INTERACTIONS

Overview

Bile acid sequestering agents may interfere with the action of URSO[®] and URSO DS[®] by reducing absorption. Aluminum based antacids adsorb bile acids *in vitro* and may act in the same manner as sequestering agents, thereby interfering with the action of URSO[®] and URSO DS[®]. Ursodiol has been shown to be an inducer of CYP3A however the clinical relevance is not known. Metabolic interactions with compounds metabolized by cytochrome P4503A are to be expected.

Drug-Drug Interactions

Table 5: Drug-Drug Interactions

	Effect	Clinical comment
Bile acid sequestrants (i.e. cholestyramine or cholestipol)	Reduces ursodiol absorption	May interfere with the action of URSO [®] and URSO DS [®]
Aluminum based antacids	Reduces ursodiol absorption Adsorbs bile acid <i>in vitro</i>	May be expected to interfere with URSO [®] and URSO DS [®]
Cytochrome P4503A substrates cyclosporine, nitrendipine and dapsone	Metabolic interaction.	Metabolic interactions with compounds metabolized by cytochrome P4503A are to be expected.

Drug-Food Interactions

Interactions with food have not been established.

Drug-Herb Interactions

Interactions with herbal products have not been established.

Drug-Laboratory Interactions

Interactions with laboratory tests have not been established.

DOSAGE AND ADMINISTRATION

Dosing Considerations

Patient Monitoring: Serum liver function tests (γ -GT, alkaline phosphatase, AST, ALT) and bilirubin level should be monitored every month for three months after start of therapy, and every six months thereafter. Serial monitoring will allow for the early detection of a possible deterioration of the hepatic function. Serum levels of these parameters usually decrease rapidly. Improved serum liver tests (e.g. AST, ALT) do not always correlate with improved disease status. For patients who have a recent history of adequate biochemical response to the treatment, UDCA discontinuation should be considered when serum liver function tests increase to a level considered clinically significant (see INDICATIONS and CLINICAL USE and WARNINGS AND PRECAUTIONS).

Caution has to be exercised to maintain the bile flow of the patients taking UDCA .

Recommended Dose

The recommended adult dosage for URSO[®] and URSO DS[®] (ursodiol) in the treatment of PBC is 13-15 mg/kg/day administered in two to four divided doses with food. The URSO DS[®] scored tablet can be broken in halves to provide recommended dosage.

Missed Dose

If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose, skip the dose you missed and take your next regularly scheduled dose. Do not take a double dose.

OVERDOSAGE

Accidental or intentional overdose with ursodiol has not been reported. The most severe manifestation of overdose would likely consist of diarrhea that should be treated symptomatically.

Symptoms of acute toxicity in animal studies were salivation and vomiting in dogs, and ataxia, dyspnea, ptosis, agonal convulsions and coma in hamsters.

For management of a suspected drug overdose, contact your regional Poison Control Centre.

ACTION AND CLINICAL PHARMACOLOGY

Mechanism of Action

Ursodiol, a naturally occurring hydrophilic bile acid, derived from cholesterol, is present as a minor fraction of the total human bile acid pool. Oral administration of ursodiol increases this fraction in a dose related manner, to become the major biliary acid, replacing/displacing toxic concentrations of endogenous hydrophobic bile acids that tend to accumulate in cholestatic liver disease.

Multiple mechanisms of action at the cellular and molecular level in addition to the replacement and displacement of toxic bile acids include cytoprotection of the injured bile duct epithelial cells (cholangiocytes) against toxic effects of bile acids, inhibition of apoptosis of hepatocytes, immunomodulatory effects via a number of mechanisms including decreasing expression of MHC class I proteins on hepatocytes and cholangiocytes, and stimulation of bile secretion by hepatocytes and cholangiocytes.

The cholesterol-lowering effect observed following the administration of URSO[®] and URSO DS[®] in patients with primary biliary cirrhosis could be related to an improvement of cholestasis, modifications in cholesterol metabolism, or both. Changes in the endogenous bile acid composition induced by URSO[®] and URSO DS[®] might be the common denominator of these two mechanisms.

Pharmacodynamics

During chronic administration, ursodiol becomes a major biliary and plasma bile acid. At a chronic dose of 13-15 mg/kg/day, ursodiol constitutes 30-50% of biliary and plasma bile acids.

Pharmacokinetics

Absorption: Ursodiol (UDCA) is normally present as a minor fraction of the total bile acids in humans (about 5%). Following oral administration, the majority of ursodiol is absorbed by passive diffusion and its absorption is incomplete.

Distribution: In healthy subjects, at least 70% of ursodiol (unconjugated) is bound to plasma protein. No information is available on the binding of conjugated ursodiol to plasma protein in

healthy subjects or primary biliary cirrhosis (PBC) patients. However, since the efficacy of ursodiol is related to its concentration in bile rather than in plasma, serum levels are not indicative of bioavailability in clinical settings. Its volume of distribution has not been determined, but is expected to be small since the drug is mostly distributed in the bile and small intestine. In bile, UDCA concentration reaches a peak in 1-3 hours.

Metabolism: Once absorbed, ursodiol undergoes hepatic extraction to the extent of about 70% in the absence of liver disease. This leads to low blood levels in the systemic circulation. As the severity of liver disease increases, the extent of extraction decreases. In the liver, ursodiol is conjugated with glycine or taurine, then secreted into bile. These conjugates of ursodiol are absorbed in the small intestine by passive and active mechanisms. The conjugates can also be deconjugated in the ileum by intestinal enzymes, leading to the formation of free ursodiol that can be reabsorbed and reconstituted in the liver. Nonabsorbed ursodiol passes into the colon where it is mostly 7-dehydroxylated to lithocholic acid. Some ursodiol is epimerized to chenodiol (CDCA) via a 7-oxo intermediate. Chenodiol also undergoes 7-dehydroxylation to form lithocholic acid. These metabolites are poorly soluble and excreted in the feces. A small portion of lithocholic acid is reabsorbed, conjugated in the liver with glycine or taurine, and sulfated at the 3 position. The resulting sulfated lithocholic acid conjugates are excreted in bile and then lost in feces.

Lithocholic acid, when administered chronically to animals, causes cholestatic liver injury that may lead to death from liver failure in certain species unable to form sulfate conjugates. Ursodiol is 7-dehydroxylated more slowly than chenodiol. For equimolar doses of ursodiol and chenodiol, steady state levels of lithocholic acid in biliary bile acids are lower during ursodiol administration than with chenodiol administration. Humans and chimpanzees can sulfate lithocholic acid. Although liver injury has not been associated with ursodiol therapy, a reduced capacity to sulfate may exist in some individuals. Nonetheless, such a deficiency has not yet been clearly demonstrated and must be extremely rare, given the several thousand patient-years of clinical experience with ursodiol.

Excretion: Ursodiol is excreted primarily in the feces. With treatment, urinary excretion increases, but remains less than 1%, except in severe cholestatic liver disease.

STORAGE AND STABILITY

URSO[®] and URSO DS[®] tablets should be stored at controlled room temperature at 15°C - 30°C in a closed container.

Half-tablets (scored URSO DS[®] tablets broken in half) maintain acceptable quality for up to 28 days when stored in the current packaging (bottles) at 15°C-30 °C .

SPECIAL HANDLING INSTRUCTIONS

To prevent leakage of the active medicinal ingredient (ursodiol) from the cut tablets onto the surface of whole tablets and spreading the bitter taste, it is recommended that cut tablets be stored separately from whole tablets

DOSAGE FORMS, COMPOSITION AND PACKAGING

250 mg tablet

URSO[®] (ursodiol) is available in a strength of 250 mg as a white, elliptical, biconvex, film-coated tablet, engraved with "URS785" on one side. Available as bottles of 100 tablets.

500 mg tablet

URSO DS[®] (ursodiol) is available in a strength of 500 mg as a white, elliptical, biconvex, scored, film-coated tablet, engraved with "URS790" on one side. Available as bottles of 100 tablets.

In addition to ursodiol as the active ingredient, URSO[®] and URSO DS[®] contain the following excipients: carnauba wax, dibutyl sebacate, ethylcellulose aqueous (cetyl alcohol, ethylcellulose, hydrogen peroxide, sodium lauryl sulfate), hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, and sodium starch glycolate.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name: Ursodiol

Chemical name: 3 α , 7 β -dihydroxy-5 β -cholan-24-oic acid

Molecular formula and molecular mass: C₂₄H₄₀O₄ 392.6

Structural formula:



Physicochemical properties:

Description: Ursodiol is a naturally occurring bile acid in man. Ursodiol is a bitter-tasting, white or almost white, crystalline powder.

Solubility: Ursodiol is practically insoluble in water, freely soluble in alcohol and glacial acetic acid, sparingly soluble in chloroform, and very slightly soluble in ether.

Melting Range: 200°C - 205°C

pKa: 6.0

pH: Alkaline

CLINICAL TRIALS

Study demographics and trial design:

Table 6: Summary of patient demographics for clinical trials in primary biliary cirrhosis (PBC)

Study	Trial design	UDCA ¹ Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range) (years)	Gender
US Study	multicenter, randomized, double-blind, placebo-controlled	13-15 mg/kg/day, administered in 4 divided doses (n=89), or placebo (n=91), 2 years	180 patients with PBC	UCDA: 53.6 (±9.5) placebo: 51.5 (±9.3)	UCDA: 7 M/89 F placebo: 12 M/91 F
Canadian Study	randomized, double-blind, placebo controlled	14 mg/kg/day (n=111), or placebo (n= 111), 2 years	222 patients with PBC	UCDA: 57.3 placebo: 55.4	UCDA: 10 M/111 F placebo: 6 M/111 F
Multinational Study	multicenter, multinational (France-Canada), double-blind, placebo controlled	13-15 mg/kg/day (n=72), or placebo (n=73), 2 years	145 patients with histologically confirmed biliary cirrhosis	UCDA: 55 (±1) placebo: 57 (±1)	UCDA: 4 M/72 F placebo: 8 M/73 F

¹ UDCA = Ursodeoxycholic acid = Ursodiol = URSO[®]

U.S. Study: A multicenter, randomized, double-blind, placebo-controlled study was conducted to evaluate the efficacy of ursodeoxycholic acid at a dose of 13-15 mg/kg/day, administered in 4 divided doses in 180 patients with PBC. Upon completion of the double-blind portion, all patients entered an open-label, active treatment, extension phase.

Treatment failure, the main efficacy end point measured during this study, was defined as death, need for liver transplantation, histologic progression by two stages or to cirrhosis, development of varices, ascites or encephalopathy, marked worsening of fatigue or pruritus, inability to tolerate the drug, doubling of serum bilirubin and voluntary withdrawal. After two years of double-blind treatment, the incidence of treatment failure was significantly reduced in the URSO[®] group (n=89) as compared to the placebo group (n=91). Time to treatment failure was also significantly delayed in the URSO[®] treated group, regardless of either histologic stage or baseline bilirubin levels (>1.8 or ≤1.8 mg/dL).

Using a definition of treatment failure which excluded doubling of serum bilirubin and voluntary withdrawal, time to treatment failure was significantly delayed in the URSO[®] group. In comparison with placebo, treatment with URSO[®] resulted in a significant improvement in the following serum hepatic biochemistries when compared to baseline: total bilirubin, AST, alkaline phosphatase and IgM.

Canadian Study: A second study conducted in Canada randomized 222 PBC patients to ursodiol 14 mg/kg/day (n=111) or placebo (n=111), in a double-blind manner during a two-year period. At

two years, a statistically significant difference between the two treatments, in favor of ursodiol, was demonstrated by the following: reduction in the proportion of patients exhibiting a more than 50% increase in serum bilirubin; median percent decrease in bilirubin, transaminases and alkaline phosphatase, incidence of treatment failure, and time to treatment failure. The definition of treatment failure included: discontinuing the study for any reason, a total serum bilirubin level greater than or equal to 1.5 mg/dL or increasing to a level equal to or greater than two times the baseline level, and the development of ascites or encephalopathy.

Evaluation of patients at 4 years or longer was inadequate due to the high dropout rate and small number of patients. Therefore, death, need for liver transplantation, histological progression by two stages or to cirrhosis, development of varices, ascites or encephalopathy, marked worsening of fatigue or pruritus, inability to tolerate the drug, doubling of serum bilirubin and voluntary withdrawal were not assessed.

Multinational Study: A two-year multicenter, multinational (France-Canada), double-blind study was conducted to compare the efficacy of ursodiol versus placebo in primary biliary cirrhosis. Patients with histologically confirmed biliary cirrhosis were randomized to receive either ursodiol 13-15 mg/kg/day (n=72), or placebo (n=73). Treatment failure was defined as a doubling of bilirubin levels (>70 µmol/L) or the occurrence of severe complications (ascites or variceal bleeding) or an adverse event.

The results showed that treatment failed in six patients in the ursodiol group, as compared with thirteen in the placebo group (p<0.01). A single patient in each group withdrew because of minor adverse effects. After two years of treatment, the proportion of patients with clinically overt disease decreased only in the ursodiol group (p<0.02). The patients treated with ursodiol had significant improvements in serum levels of bilirubin, alkaline phosphatase, alanine aminotransferase, aspartate aminotransferase, γ-glutamyltransferase, cholesterol, and IgM (all p<0.01); the antimitochondrial antibody titer (p<0.01); and the Mayo risk score (p<0.001). In a follow-up analysis of 95 liver-biopsy specimens, only the group receiving ursodiol showed a significant improvement in the mean histologic score (p<0.002) and in all the characteristic histologic features except fibrosis.

At the end of this trial, all patients received ursodiol (13-15 mg/kg/day) and were monitored for an additional two years, using the same criteria.

After four years, the overall treatment failure rate was 12% in the ursodiol group and 26% in the original placebo group (p<0.001). Two patients in the ursodiol group had undergone a liver transplantation, compared to 12 in the original placebo group (p<0.001). Survival was similar in the two groups: 5 deaths (various causes) occurred in the ursodiol group and 7 in the original placebo group.

Combined Analysis: The raw data from the above three studies have been combined in order to estimate, at four years, the magnitude of the effect of ursodiol treatment on survival free of transplant, defined as time to transplant, or death without transplant.

In these studies, all patients had histologically confirmed, antimitochondrial antibody positive, primary biliary cirrhosis. They were randomized to receive ursodiol (13-15 mg/kg/day) or identical

placebo. In one study, blinded randomization continued for four years. In two studies, open label ursodiol was offered to all patients after two years. The endpoint of survival free of liver transplant was compared between the ursodiol and placebo groups using standard life table analyses. Analyses were done on an "intent-to-treat" basis. The risk reduction was calculated in order to define the magnitude of the benefit from ursodiol treatment.

A total of 548 patients were randomized in these studies: 273 received ursodiol and 275, the placebo. Baseline characteristics were comparable at entry in both groups. Median length of follow-up was four years in the ursodiol group, and 3.8 years in the original placebo group. Placebo patients who received ursodiol did so for a mean of one year. There were 47 patients in the ursodiol group and 68 in the placebo group who did not survive nor needed a liver transplant. Survival free of transplantation was extended in patients originally randomized to ursodiol when compared to those on placebo (mean of 3.66 versus 3.45 years, $p=0.014$). In the ursodiol group the risk of dying or being transplanted was reduced by 32% ($\pm 11\%$) of that observed in the original placebo group.

Unapproved High-Dose Ursodeoxycholic Acid for the Treatment of Primary Sclerosing Cholangitis: In a recent Clinical Trial (Lindor *et al.*, 2009), one hundred fifty adult patients with PSC were enrolled in a long-term, randomized, double-blind controlled trial of high-dose (28-30 mg/kg/day – 1.5 to 2.0 fold the recommended dose) versus placebo. Liver biopsy and cholangiography were performed before randomization and after 5 years. The primary outcome measures were development of cirrhosis, varices, cholangiocarcinoma, liver transplantation, or death. The study was terminated after 6 years due to futility. During therapy, aspartate aminotransferase and alkaline phosphatase levels decreased more in the ursodeoxycholic acid group than the placebo group ($P<0.01$), but improvements in liver function tests were not associated with decreased endpoints. By the end of the study, 30 patients in the ursodeoxycholic acid group (39%) versus 19 patients in the placebo group (26%) had reached one of the pre-established clinical endpoints. The risk was 2.1 times greater for death and transplantation in the ursodeoxycholic acid group versus the placebo group ($P = 0.038$). Serious adverse events were more common in the ursodeoxycholic acid group than the placebo group (63% versus 37% [$P < 0.01$]). Long-term, high-dose ursodeoxycholic acid therapy was associated with improvement in serum liver function tests in PSC but did not improve survival and was associated with higher rates of serious adverse events.

DETAILED PHARMACOLOGY

Administration of ursodiol to rats, rabbits, hamsters and dogs produced modification of bile composition. Bile flow increased as did total bile acid output. In the liver, ursodiol decreased HMG-CoA reductase activity and cholesterol 7-hydroxylase activity. Triglyceride, phospholipid and cholesterol synthesis were decreased.

Studies have demonstrated that ursodiol acts on the hepatic cells and plays a role in the bile acid dependent mechanism of bile formation. Its choleric activity results from its osmotic activity as well as its stimulating effect on organic ion transport (probably HCO_3).

In vitro studies showed that tauroursodeoxycholic acid (i.e. in the liver ursodiol is conjugated with taurine or glycine) decreased cholesterol uptake in rat jejunal membranes by an unknown mechanism. When ursodiol was perfused into the liver of rats or baboons, bile flow either remained unchanged or increased, bile acid and phospholipid outputs were increased, while cholesterol specific activity was decreased. Tauroursodeoxycholic acid caused only little output of plasma membrane enzyme concentration (5-nucleotidase and alkaline phosphatase), which may represent a characteristic difference between the effects of chenodiol and ursodiol on the hepatobiliary system.

Ursodiol produced minimal or no effect on water and sodium excretion from the GI tract of rats and rabbits. It induced less damage to the GI tract mucosa than chenodiol. These observations correlate well with the clinical findings that diarrhea is infrequent with ursodiol.

Ursodiol lowered blood sugar levels in mice, and increased the volume of pancreatic secretion in rabbits, thus suggesting a stimulatory effect of ursodiol on the pancreas.

At therapeutic doses, ursodiol uncouples the normal relationship between cholesterol, phospholipids and bile acid secretion. Ursodiol inhibits cholesterol absorption in the gut, thereby, reducing cholesterol output into the bile. It further reduces cholesterol secretion into bile. These actions contribute to biliary cholesterol desaturation.

TOXICOLOGY

Acute Toxicity

Results from various studies indicated that oral, subcutaneous, intraperitoneal and intravenous administration of ursodiol in mice, rats, hamsters, and dogs at single doses of 1.21 to 10 g/kg over a seven-day observation period, did not cause any deaths in any of the species used. For mice and dogs, the LD₅₀ was >10 g/kg, and rats had an LD₅₀ over >5 g/kg. Hamsters were found to be more sensitive than rats and dogs as the LD₅₀ for this species was calculated to be >3.16 g/kg.

No significant sex difference was seen. Toxic signs observed included: inhibition of motility, CNS toxicity such as ataxia and sedation, GI tract disturbances such as vomiting, salivation, decreased body weight and appetite.

Subacute Toxicity

Two short-term toxicological studies were conducted in rats. Ursodiol was administered orally at a daily dose of 0.5 to 4.0 g/kg/day for five weeks or alternatively at doses of 0.0625 to 0.5 g/kg daily for five weeks by the intraperitoneal route.

No deaths occurred in the study with oral administration of ursodiol, whereas, one male and one female rat died in the 0.25 g/kg group, and six males and four females died in the 0.5 g/kg group of the study in which ursodiol was administered by the intraperitoneal route. The most marked autopsy findings were dilation and adhesion of intraperitoneal organs. As these became gradually more severe, retention of ascites and renal abscesses appeared. It was concluded that 0.0625 g/kg was the safe dose and 0.125 g/kg was near the maximum tolerable dose.

Ursodiol orally administered to rats did not cause any clinical symptoms or any changes in laboratory parameters.

Chronic Toxicity

Four long-term toxicity studies were performed in rats and monkeys. The results of these studies are summarized below.

Rat Study: In one study, ursodiol was administered orally to Sprague-Dawley rats for 26 weeks. The dosage varied between 0.1 and 2.5 g/kg/day and various observations were performed daily.

No deaths occurred during the experimental period. Lower doses (0.1 and 0.5 g/kg) were well tolerated. However, a 2.5 g/kg dose of ursodiol resulted in significant reduction of body weight gain and food intake. No significant changes were seen in laboratory findings and clinical observations.

In the second study, male Wistar rats were given 0.5 to 4.0 g/kg of ursodiol orally for 26 consecutive weeks and a variety of observations were made.

The results indicated a decrease in body weight gain and an increase in water intake in the 4.0 g/kg dosage group. Eight rats (four at the high dose level) died during the experiment. The cause of death was attributed to pathological changes in the lung and intestine. Laboratory findings revealed no abnormal changes that might be ascribed to drug administration.

Monkey study: A 26-week study was performed in Rhesus monkeys. Ursodiol at doses of 0.04 and 0.10 g/kg/day were given orally.

No deaths occurred during the treatment period. There were no abnormalities in the laboratory parameters.

In a 52-week study, ursodiol at a dose of 0.05 to 0.9 g/kg was administered to Rhesus monkeys. The animals were observed daily for various clinical signs and symptoms. They were weighed weekly, blood and urine was collected and examined every three months. After 52 weeks, the animals were sacrificed and an autopsy was performed.

Three animals belonging to the 0.90 g/kg group, two in the 0.30 g/kg group and one in the 0.10 g/kg died during the study. These deaths were considered to be related to ursodiol. Liver toxicity (small round-cell infiltration, vacuolar degeneration, necrosis of hepatic cells, phagocytosis and hepatic abscess) and thickening of the alveolar wall of the lungs was observed in deceased animals from all groups. Necrosis of the stomach wall was observed in deceased animals from the 0.90 g/kg group. A regression of body weight gain was seen in the 0.30 and 0.90 g/kg groups. Episodes of diarrhea were observed in all groups including the control group. No remarkable changes were noted in hematological, urinary, electrographic, blood pressure and ocular fundi examinations. However, serum SGPT, AST and ALP increased significantly.

From the above findings, it was concluded that ursodiol, when administered at daily doses exceeding 0.10 g/kg, caused hepatotoxicity in Rhesus monkeys.

Carcinogenicity

In two 24-month oral carcinogenicity studies in mice, ursodiol at doses up to 1,000 mg/kg/day (3,000 mg/m²/day) was not tumorigenic. Based on body surface area, for a 50 kg person of average height (1.46 m² body surface area), this dose represents 5.4 times the recommended maximum clinical dose of 15 mg/kg/day (555 mg/m²/day).

In a two-year oral carcinogenicity study in Fischer 344 rats, ursodiol at doses up to 300 mg/kg/day (1,800 mg/m²/day, 3.2 times the recommended maximum human dose based on body surface area) was not tumorigenic.

In a life-span (126-138 weeks) oral carcinogenicity study, Sprague-Dawley rats were treated with doses of 33 to 300 mg/kg/day, 0.4 to 3.2 times the recommended maximum human dose based on body surface area. Ursodiol produced a significantly ($p < 0.5$, Fisher's exact test) increased incidence of pheochromocytomas of the adrenal medulla in females of the highest dose group.

In 103-week oral carcinogenicity studies of lithocholic acid, a metabolite of ursodiol, doses up to 250 mg/kg/day in mice and 500 mg/kg/day in rats did not produce any tumors. In a 78-week rat study, intrarectal instillation of lithocholic acid (1 mg/kg/day) for 13 months did not produce colorectal tumors. A tumor-promoting effect was observed when it was administered after a single intrarectal dose of a known carcinogen N-methyl-N'-nitro-N-nitrosoguanidine. On the other hand, in a 32-week rat study, ursodiol at a daily dose of 240 mg/kg (1,440 mg/m², 2.6 times the maximum recommended human dose based on body surface area) suppressed the colonic carcinogenic effect of another known carcinogen, azoxymethane.

Mutagenicity

Ursodiol was not genotoxic in the Ames test, the mouse lymphoma cell (L5178Y, TK^{+/-}) forward mutation test, the human lymphocyte sister chromatid exchange test, the mouse spermatogonia chromosome aberration test, the Chinese hamster micronucleus test and the Chinese hamster bone marrow cell chromosome aberration test.

Reproduction and Teratology

Ursodiol did not show any teratogenic effect in mice, rats and rabbits at oral dose levels up to 1.5, 4.0 and 0.3 g/kg, respectively, and in mice and rats at intraperitoneal dose levels up to 0.2 g/kg. Furthermore, it did not influence mating performance and fertility, except in one study where these parameters were slightly reduced in female rats receiving 2.0 g/kg. Breeding capacity was not altered by the administration of ursodiol.

Oral administration of 1.5 g/kg in mice and 2.0 g/kg in rats induced a decrease in maternal weight gain and lower mean weights of live fetuses. In addition, the number of resorption sites was increased in rats at a dose of 2.0 g/kg. Rabbits were much more sensitive than mice and rats to the toxic action of ursodiol. The administration of doses of 0.1 g/kg and greater caused a decrease in food consumption, maternal body weight gain and motor activity as well as an increase in resorption sites, and absorption death.

Intraperitoneal administration of 0.2 g/kg ursodiol to mice and rats induced a decrease in maternal body weight gain, low fetal weight and an increase of resorption sites.

REFERENCES

1. Dubner H, Fromm H. Ursodeoxycholic acid treatment of intrahepatic cholestasis of pregnancy: observations on efficacy and safety. *Gastroenterology* 1993; 104(2):660-661.
2. Mazzella G, Rizzo N, Azzaroli F, Simoni P, Bovicelli L, Miracolo A, Simonazzi G, Colecchia A, Nigro G, Mwangemi C, Festi D, Roda E. Ursodeoxycholic acid administration in patients with cholestasis of pregnancy: effects on primary bile acids in babies and mothers. *Hepatology* 2001; 33(3):504-508.
3. Roberts IM. Hyperlipidemic gestational pancreatitis. *Gastroenterology* 1993; 104(5): 1560-1562.
4. Poupon R. Primary biliary cirrhosis: immunological aggression and liver injury from bile acids. *HRLR* 1992; 9: XI-XIV.
5. Lindor KD, Dickson ER, Baldus WP, Jorgensen RA, Ludwig J, Murtaugh PA, Harrison JM, Wiesner RH, Anderson ML, Lange SM, Lesage G, Rossi SS, Hofmann AF. Ursodeoxycholic acid in the treatment of primary biliary cirrhosis. *Gastroenterology* 1994; 106(5):1284-1290.
6. Clinical Study Report: A cross-over trial evaluating single- and multiple- daily dosage schedules of ursodeoxycholic acid in the treatment of primary biliary cirrhosis. Data on file. Axcan Pharma Inc.
7. Hempfling W, Dilger K, Beuers U. Systemic review: ursodeoxycholic acid-adverse effects and drug interactions. *Aliment Pharmacol Ther.* 2003; 18(10): 963-972.
8. Angulo P, Dickson ER, Therneau TM, Jorgensen RA, Smith C, DeSotel CK, Lange SM, Anderson ML, Mahoney DW, Lindor KD. Comparison of three doses of ursodeoxycholic acid in the treatment of primary biliary cirrhosis: a randomized trial. *Hepatology* 1999; 30:830-835.
9. Angulo P, Lindor KD. Management of primary biliary cirrhosis and autoimmune cholangitis. *Clin Liver Dis* 1998; 2(2):333-351.
10. Fedorowski T, Salen G, Colallilo A, Tint GS, Mosbach EH, Hall JC. Metabolism of ursodeoxycholic acid in man. *Gastroenterology* 1977; 73(5):1131-1137.
11. Poupon RE, Balkau B, Eschwège E, Poupon R. A multicenter, controlled trial of ursodiol for the treatment of primary biliary cirrhosis. *N Engl J Med* 1991; 324:1548-1554.
12. Poupon R, Calmus Y, Poupon RE. Ursodeoxycholic acid for primary biliary cirrhosis: effects and mechanisms of action. *Cellular and Molecular Aspects of Cirrhosis.* 1992; 216: 345-349.

13. Podda M, Ghezzi C, Battezzati PM, Bertolini E, Crosignani A, Petroni ML, Zuin M. Ursodeoxycholic acid for chronic liver diseases. *J Clin Gastroenterol* 1988; 10(Suppl 2):S25-S31.
14. Angulo P. Use of ursodeoxycholic acid in patients with liver disease. *Curr Gastroenterol Rep* 2002; 4:37-44.
15. Trauner M, Graziadei IW. Review article: mechanisms of action and therapeutic applications of ursodeoxycholic acid in chronic liver disease. *Aliment Pharmacol Ther.* 1999; 13:979-995.
16. Kumar D, Tandon RK. Use of ursodeoxycholic acid in liver diseases. *J Gastroenterol and Hepatol* 2001; 16:3-14.
17. Kowdley KV. Ursodeoxycholic acid thereapy in hepatobiliary disease. *Am J Med* 2000; 108:481-486.
18. Paumgartner G, Beuers U. Mechanisms of action and therapeutic efficacy of ursodeoxycholic acid in cholestatic liver disease. *Clin Liver Dis.* 2004; 8(1):67-81.
19. Mazzella G, Parini P, Festi D, Bazzoli F, Aldini R, Roda A, Tonelli D, Cipolla A, Salzetta A, Roda E. Effect of simvastatin, ursodeoxycholic acid and simvastatin plus ursodeoxycholic acid on biliary lipid secretion and cholic acid kinetics in nonfamilial hypercholesterolemia. *Hepatology* 1992; 15(6): 1072-1078.
20. Bazzoli F, Marselli Labate AM, Malavolti M, Roda A, Monti F, Rucci P, Roda E. The effect of ursodeoxycholic administration on bile lipid composition in man: Comparison of three different ursodeoxycholic preparations. *Curr Ther Res* 1983; 33(3):543-550.
21. Poupon RE, Lindor KD, Cauch-Dudek K, Dickson ER, Poupon R, Heathcote EJ. Combined analysis of randomized controlled trials of ursodeoxycholic acid in primary biliary cirrhosis. *Gastroenterology* 1997; 113(3):884-890.
22. Matern S, Tietjen KG, Fackler O, Hinger K, Herz R, Gerok W. Bioavailability of ursodeoxycholic acid in man: Studies with a radioimmunoassay for ursodeoxycholic acid. In: G. Paumgartner, A. Stiechl, W. Gerole (eds.): *Biological Effects of Bile Acids*, Falk Symp. No. 26. MTP Press Limited, Lancaster (1979): 109-118.
23. Hoshita T, Kono M, Matsumoto M, Uchiyama M, Kuramoto T. Metabolism of bile acids. I. Absorption, distribution, excretion and metabolism of ursodeoxycholic acid. *J Pharm Soc Japan* 1974; 94:1196-1205.

24. Siegel JL, Jorgensen R, Angulo P, Lindor KD. Treatment With Ursodeoxycholic Acid Is Associated With Weight Gain in Patients With Primary Biliary Cirrhosis. *J Clin Gastroenterol* 2003; 37(2):183-185.
25. Stefaniwsky AB, Tint GS, Speck J, Salen G. Ursodeoxycholic acid (UDCA) reduces pain, nausea, and vomiting in patients with bile acid reflux gastritis. *Gastroenterology* 1982; 82:1188.
26. Heathcote EJ, Cauch-Dudek K, Walker V, Baily RJ, Blendis LM, Ghent CN, Michieletti P, Minuk GY, Pappas SC, Scully LJ, Steinbrecher UP, Sutherland LR, Williams CN, Witt-Sullivan H, Worobetz LJ, Milner RA, Wanless IR. The Canadian multicenter double-blind randomised controlled trial of ursodeoxycholic acid in primary biliary cirrhosis. *Hepatology* 1994; 19(5):1149-1156.
27. Poupon RE, Poupon R and Balkau B. Ursodiol for the long-term treatment of primary biliary cirrhosis. *N Engl J Med* 1994; 330(19):1342-1347.
28. Angulo P, Batts KP, Therneau TM, Jorgensen RA, Dickson, ER Dickson, Lindor KD. Long-term ursodeoxycholic acid delays histological progression in primary biliary cirrhosis. *Hepatology* 1999; 29(3):644-647.
29. Bellentani S, Manenti F, and Ferrari A. Meta-analysis of the clinical results with UDCA in chronic liver diseases. In: Falk symposium 71: Fat-storing cells and liver fibrosis. Surrenti C, Casini A, Milani S, Pinzani M (eds). Kluwer Academic Publishers. Dordrecht, Boston, London 1994: 340-347.
30. Corpechot C, Carrat F, Bonnard AM, Poupon RE, Poupon R. The effect of ursodeoxycholic acid therapy on liver fibrosis progression in primary biliary cirrhosis. *Hepatology* 2000; 32(6):1196-1199.
31. Degott C, Zafrani ES, Callard P, Balkau B, Poupon RE, Poupon R. Histopathological study of primary biliary cirrhosis and the effect of ursodeoxycholic acid treatment on histology progression. *Hepatology* 1999; 29(4):1007-1012.
32. Jorgensen R, Angulo P, Dickson ER, Lindor KD. Results of long-term ursodiol treatment for patients with primary biliary cirrhosis. *Am J Gastroenterol.* 2002; 97(10):2647-2650.
33. Lindor KD, Therneau TM, Jargensen RA, Malinchoc M, Dickson ER. Effects of ursodeoxycholic acid on survival in patients with primary biliary cirrhosis. *Gastroenterology* 1996; 110(5):1515-1518.
34. Ludwig J, Dickson ER, McDonald GS. Staging of chronic nonsuppurative destructive cholangitis (syndrome of primary biliary cirrhosis). *Virchows Archiv A Path Anat and Histo* 1978; 379: 103-112.
35. Poupon RE, Bonnard AM, Chrétien Y, Poupon R. Ten-year survival in ursodeoxycholic

- acid treated patients with primary biliary cirrhosis. *Hepatology* 1999; 29(6):1668-1671.
36. Simko V, Michael S, Prego V. Ursodeoxycholic therapy in chronic liver disease: a meta-analysis in primary biliary cirrhosis and in chronic hepatitis. *Am J Gastroenterol* 1994; 89(3):392-398.
 37. Lee D, Bonorris G, Cohen H, Gilmore C, Marks J, Schoenfield LJ. Effect of ursodeoxycholic acid on bile acid kinetics and hepatic lipid secretion. *Hepatology* 1981; 1:36A.
 38. Logan GM and Duane WC. Lovastatin added to ursodeoxycholic acid further reduces biliary cholesterol saturation. *Gastroenterology* 1990; 98(6):1572-1576.
 39. Takahashi H, Tozuka K, Miyashita T, et al. Effects of ursodeoxycholic acid administered to primipara rats and mice on the prenatal development and postnatal growth of their litters. *Clinical Report* 1975; 9(13): 3223-3242.
 40. Toyoshima S, Fujita H, Sato R, Kashima M, Sato S. Reproduction studies with ursodeoxycholic acid in rats. (I) fertility study. *Pharmacometrics* 1978; 15(5): 923-930.
 41. Toyoshima S, Fujita H, Sakurai T, Sato R, Kashima M. Reproduction studies with ursodeoxycholic acid in rats. (II) teratogenicity study. *Pharmacometrics* 1978; 15(5): 931-945.
 42. Toyoshima S, Fujita H, Sato R, Kashima M, Sato S. Terotogenicity study of ursodeoxycholic acid in rabbits. *Pharmacometrics* 1978; 5(7):1133-1140.
 43. Lindor *et al.* High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology* 2009; 50(3): 808-814.

PART III: CONSUMER INFORMATION

PrURSO®
250 mg tablets

PrURSO DS®
500 mg tablets

ursodiol tablets

This leaflet is part III of a three-part "Product Monograph" and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about URSO® and URSO DS®. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS MEDICATION

What the medication is used for:

URSO® or URSO DS® has been prescribed for you by your physician for the management of cholestatic liver disease.

URSO® and URSO DS® are only available by prescription.

What it does:

URSO® and URSO DS® contain ursodiol (ER-so-DYE-all), which is a naturally occurring bile acid found in small quantities in normal human bile.

In patients with cholestatic liver disease the release and flow of bile through the bile ducts are reduced. By taking URSO® or URSO DS® the amount of ursodiol in the bile increases, changing the make-up of the bile and causing an increase in bile flow. Ursodiol also works by replacing toxic bile acids that can destroy liver tissue.

When it should not be used:

Do not take URSO® or URSO DS® if you:

- have an allergy to ursodiol or to any ingredient in the formulation.
- have blockage of bile flow due to widespread liver disease.
- have complete blockage of bile flow due to disease outside of the liver.

What the medicinal ingredient is:

ursodiol

What the nonmedicinal ingredients are:

URSO® and URSO DS® contain the following excipients: carnauba wax, dibutyl sebacate,

ethylcellulose aqueous (cetyl alcohol, ethylcellulose, hydrogen peroxide, sodium lauryl sulfate), hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, and sodium starch glycolate.

What dosage forms it comes in:

250 mg tablet

URSO® (ursodiol) is available as 250 mg white, elliptical, biconvex, film-coated tablets, engraved with "URS785". Each bottle contains 100 tablets.

500 mg tablet

URSO DS® (ursodiol) is available as 500 mg white, elliptical, biconvex, scored, film-coated tablets, engraved with "URS790". Each bottle contains 100 tablets.

WARNINGS AND PRECAUTIONS

Before you take URSO® or URSO DS®, tell your doctor if:

- You have taken URSO® or URSO DS® before and it was not well-tolerated or caused an allergy.
- You have liver problems, or are in need of a liver transplant.
- You have a partial blockage of bile flow due to disease outside of the liver.
- You have variceal bleeding (bleeding from swollen veins, arteries, or lymph vessels).
- You have ascites (swelling in the abdomen).
- You are taking other prescription or non-prescription medicines.
- You are using any new medicine (prescription or non-prescription) such as bile reduction medicines (cholestyramine or colestipol), aluminum based antacids (Rolaids, Maalox, Mylanta, and many others), and cyclosporine, nifedipine, or dapsone.
- You develop any new medical problem while using this medicine.
- You have severe stomach pain.

- You are pregnant, plan to become pregnant, are breast-feeding or plan to breast-feed.
- You need other medical treatment by another doctor, let him or her know that you are taking URSO[®] or URSO DS[®].

URSO[®] and URSO DS[®] are not recommended for use in children.

You should discuss with your doctor the benefits and risks of taking URSO[®] and URSO DS[®] for your medical condition.

This medication is prescribed for a particular health problem and for your personal use only. Do not give it to another person.

Keep this and all other medicines out of the reach of children.

INTERACTIONS WITH THIS MEDICATION

The following medicines may decrease the amount of ursodiol that is absorbed into your body:

- Medicines that reduce the amount of bile acids such as cholestyramine or colestipol
- Antacids that contain aluminum such as Rolaids, Maalox, Mylanta, and many others.

The absorption and metabolism of the following medicines may be affected by taking ursodiol:

- Cyclosporine
- Dapsone
- Nitrendipine

Use of these medicines with URSO[®] or URSO DS[®] may require patients to be closely monitored and the dose of their medicines adjusted.

PROPER USE OF THIS MEDICATION

Usual Adult Dose:

Your doctor would have prescribed the amount of URSO[®] or URSO DS[®] you should take each day for your medical condition. URSO[®] and URSO DS[®] should be taken in 2 to 4 divided doses with food. It is easier to remember to take your medication if it is taken at the same time each day. Setting up a routine to take your medication helps this activity become a normal part of your day.

Take URSO[®] and URSO DS[®] for the full duration of

treatment, even if you begin to feel better.

To break URSO DS[®] scored tablet easily, place the tablet on a flat surface with the scored section on top. Hold the tablet with your thumbs placed close to the scored part of the tablet (groove). Then apply gentle pressure and snap the tablet segments apart (segments breaking incorrectly should not be used). The segments should be swallowed unchewed, with water.

To prevent leakage of the active medicinal ingredient (ursodiol) from the cut tablets onto the surface of whole tablets and spreading the bitter taste, it is recommended that cut tablets be stored separately from whole tablets.

This medication should only be used as instructed by your doctor. Follow your doctor's instructions. Do not change the dose or stop the treatment without your doctor's advice.

Your doctor will ask you to have regular medical checkups, and will likely require liver tests. It is important to respect the dates proposed.

Overdose:

The most severe symptom of overdosage would likely be diarrhea.

If you take more medication than you have been told to take, you should contact your doctor, hospital emergency department, or nearest poison control center immediately, even if you do not feel sick or you have no other symptoms.

Missed Dose:

If you miss a dose, take the missed dose as soon as you remember. If it is almost time for your next dose, skip the dose you missed and take your next regularly scheduled dose. Do not take a double dose.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect		Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
		Only if severe	In all cases	
Common >1% and <10%	Anemia		√	
	Dizziness	√		
	Reduced white blood cells in the blood		√	
	Headache	√		
	Diarrhea		√	
	Swelling of the extremities		√	
	Blood glucose increased		√	
	Abdominal pain		√	
Unknown	Swelling beneath the skin and Swelling of the throat			√
	Palpitation	√		
	Cough	√		
	Increase in eosinophils in the blood		√	
	Drug hypersensitivity to include facial edema			√
	Decreased blood neutrophils		√	
	Interstitial lung disease			√
	Decreased blood platelets		√	
	Severe inflammation and peeling of the skin			√
	Constipation		√	
	Skin redness and Papular skin lesion			√
	Fever		√	
	Photosensitivity reaction			√
	Aggravation or occurrence of Jaundice			√

This is not a complete list of side effects. For any unexpected effects while taking URSO® or URSO DS®, contact your doctor or pharmacist.

HOW TO STORE IT

URSO® and URSO DS® tablets should be stored at room temperature at 15°C to 30°C in a closed container.

Keep this and all other medication out of the reach of children.

REPORTING SUSPECTED SIDE EFFECTS

You can report any suspected adverse reactions associated with the use of health products to the Canada Vigilance Program by one of the following 3 ways:

- Report online at www.healthcanada.gc.ca/medeffect
- Call toll-free at 1-866-234-2345
- Complete a Canada Vigilance Reporting Form and:
 - Fax toll-free to 1-866-678-6789, or
 - Mail to: Canada Vigilance Program
Health Canada
Postal Locator 0701E
Ottawa, Ontario
K1A 0K9

Postage paid labels, Canada Vigilance Reporting Form and the adverse reaction reporting guidelines are available on the MedEffect™ Canada Web site at www.healthcanada.gc.ca/medeffect.

NOTE: Should you require information related to the management of side effects, contact your health professional. The Canada Vigilance Program does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at: <http://www.aptalispharma.com>, or by contacting the sponsor, Aptalis Pharma Canada Inc. at: 1-800-565-3255

This leaflet was prepared by Aptalis Pharma Canada Inc.

URSO® and URSO DS® are registered trademarks of Aptalis Pharma Canada Inc.

Last revised: August 18, 2014