


## GENOPTIC® Eye Drops

### NAME OF THE DRUG

The active constituent of GENOPTIC® eye drops is gentamicin sulfate.



(structure of gentamicin)

Gentamicin C <sub>1</sub> :	R <sub>1</sub> = R <sub>2</sub> = CH <sub>3</sub>
Gentamicin C <sub>1A</sub> :	R <sub>1</sub> = R <sub>2</sub> = H
Gentamicin C <sub>2</sub> :	R <sub>1</sub> = CH <sub>3</sub> R <sub>2</sub> = H

### DESCRIPTION

Gentamicin sulfate is a white to buff hygroscopic powder which is freely soluble in water, ethylene glycol and formamide. It is practically insoluble in alcohol, acetone, chloroform and ether. Gentamicin sulfate is a mixture of the sulfates of gentamicin C<sub>1</sub>, gentamicin C<sub>1A</sub> and gentamicin C<sub>2</sub>. When dry, gentamicin sulfate contains not less than 590 units of gentamicin per mg.

**Chemical Name:** O-3-deoxy-4-C-methyl-3-(methylamino)-β-L-arabinopyranosyl-(1→6)-O-[2,6-diamino-2,3,4,6-tetradeoxy-α-D-erythro-hexopyranosyl-(1→4)]-2-deoxy-D-streptamine.

**Empirical formula:**

Gentamicin C <sub>1</sub> :	C <sub>21</sub> H <sub>43</sub> N <sub>5</sub> O <sub>7</sub>
Gentamicin C <sub>1A</sub> :	C <sub>19</sub> H <sub>39</sub> N <sub>5</sub> O <sub>7</sub>
Gentamicin C <sub>2</sub> :	C <sub>20</sub> H <sub>41</sub> N <sub>5</sub> O <sub>7</sub>

GENOPTIC® eye drops are a sterile, aqueous solution buffered to approximately pH 7.0 for use in the eye. Each mL contains gentamicin sulfate (equivalent to 3 mg gentamicin) with polyvinyl alcohol (LIQUIFILM®), disodium edetate, sodium phosphate dibasic, sodium chloride, benzalkonium chloride as a preservative and purified water.

### PHARMACOLOGY

Gentamicin sulfate is a water soluble antibiotic of the aminoglycoside group which has shown activity against a wide variety of pathogenic gram-negative and gram-positive bacteria. The gram-positive bacteria against which gentamicin sulfate is active include coagulase positive and negative staphylococci.

The gram-negative bacteria against which gentamicin sulfate is active include certain strains of *Pseudomonas aeruginosa*, indole positive and indole negative *Proteus* species, *Escherichia coli*, *Klebsiella pneumoniae*, (Friedlander's bacillus), *Haemophilus influenzae*

and *Haemophilus aegyptius* (Koch-Weeks bacillus), *Aerobacter aerogenes*, *Moraxella lacunata* (diplobacillus of Morax-Axenfeld), and *Neisseria* species, including *Neisseria gonorrhoeae*.

At this time there are increasing members of resistant cases being reported for the aminoglycoside class of antibiotics, particularly to *Streptococcus pneumoniae*. This phenomenon will also reduce the synergy with the  $\beta$ -lactam class of drugs as combination therapy. This should be considered when commencing therapy for all ocular infections and the results of therapy should be closely monitored for signs of inefficiency. Should this occur, more aggressive pharmacotherapeutic options, e.g. a cephalosporin, should be used.

## PHARMACOKINETICS

Studies on the pharmacokinetics of ophthalmic preparations of gentamicin sulfate have not been conducted.

## INDICATIONS

GENOPTIC<sup>®</sup> eye drops are indicated in the topical treatment of infections of the external eye and its adnexa caused by susceptible bacteria. Such infections include conjunctivitis, keratitis, keratoconjunctivitis, corneal ulcers, blepharitis, blepharoconjunctivitis, acute meibomianitis and dacryocystitis.

## CONTRAINDICATIONS

Hypersensitivity to gentamicin sulfate, benzalkonium chloride or any of the other constituents.

## PRECAUTIONS

GENOPTIC<sup>®</sup> eye drops are not for injection. It should never be injected subconjunctivally, nor should it be introduced directly into the anterior chamber of the eye.

Prolonged use of topical antibiotics may give rise to overgrowth of nonsusceptible organisms, such as fungi. Should this occur, or if irritation or hypersensitivity to any component of the product develops, discontinue use of the preparation and institute appropriate therapy.

### **Use in pregnancy:** Category D.

Gentamicin and other aminoglycosides cross the placenta. There is evidence of selective uptake of gentamicin by the foetal kidney resulting in damage (probably reversible) to immature nephrons. Eighth cranial nerve damage has also been reported following *in-utero* exposure to some of the aminoglycosides. Because of their chemical similarity, all aminoglycosides must be considered potentially nephrotoxic and ototoxic to the foetus. It should also be noted that therapeutic blood levels in the mother do not equate with safety for the foetus.

**Use in lactation:** Studies on the use of GENOPTIC<sup>®</sup> eye drops during lactation have not been conducted.

## ADVERSE REACTIONS

Transient irritation has been reported with the use of GENOPTIC<sup>®</sup> eye drops.

## **OVERDOSAGE**

In case of overdosage, immediately flush the eye(s) with water or normal saline.

## **DOSAGE AND ADMINISTRATION**

Instil one or two drops of GENOPTIC® eye drops into the affected eye(s) every four hours. In severe infections, dosage may be increased to a maximum of 2 drops once every hour.

In order to minimise systemic absorption of GENOPTIC® eye drops, apply pressure to the tear duct immediately following administration of the drug.

## **PRESENTATION**

Eye drops: 5 mL (dropper bottle).

Storage: Store below 25°C.

Shelf life: 2 years.

To avoid contamination of the solution, keep container tightly closed.

Do not touch dropper tip to any surface.

Discard unused contents 4 weeks after opening the bottle.

Contents are sterile if seal is intact.

AUST R 23215

Allergan Australia Pty. Ltd.

810 Pacific Highway

Gordon NSW 2072

ABN: 85 000 612 831

**Poisons schedule:** S4

**TGA Approval Date:** 22 April, 1996

**Date of Most Recent Amendment:** 12 May, 2008

® Registered Trademark of Allergan Inc